



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ARITHMETIC.

107. Proposed by REBER V. ALLEN, Hooker Station, Ohio.

A barn, $ABCD$, length $AB=b$ feet, width $AD=a$ feet, standing in an open field, has a horse tethered to a point, P , in the side, AB , distance $AP=c$ feet, with a rope R feet long. Over what area can the horse graze?

I. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

We might consider six cases of which only the fifth and sixth are represented in the figure.

Case I. In every case $c < b - c$. $R > c$ and $< b - c$, also $< a + c$.

$$\text{Area} = \frac{1}{2}\pi R^2 + \frac{1}{2}\pi(R-c)^2 = \frac{1}{2}\pi(3R^2 - 2Rc + c^2).$$

Case II. $R > c$, $R > b - c$, $R < a + c$, $R < a + b - c$.

$$\text{Area} = \frac{1}{2}\pi R^2 + \frac{1}{2}\pi(R-c)^2 + \frac{1}{2}\pi(R-b+c)^2 = \frac{1}{2}\pi(4R^2 + 2c^2 + b^2 - 2Rb - 2bc).$$

Case III. $R > c + a$, $R < a + b - c$.

$$\text{Area} = \frac{1}{2}\pi R^2 + \frac{1}{2}\pi(R-c)^2 + \frac{1}{2}\pi(R-c-a)^2 + \frac{1}{2}\pi(R-b+c)^2 = \frac{1}{2}\pi(5R^2 - 2Rc + 3c^2 + a^2 - 2Ra + 2ac + b^2 - 2Rb - 2bc).$$

Case IV. $R > c + a$, $R > a + b - c$, $R < a + b$.

$$\text{Area} = \frac{1}{2}\pi R^2 + \frac{1}{2}\pi(R-c)^2 + \frac{1}{2}\pi(R-c-a)^2 + \frac{1}{2}\pi(R-b+c)^2 + \frac{1}{2}\pi(R-a-b+c)^2 = \frac{1}{2}\pi(3R^2 + 2c^2 + a^2 - Ra + b^2 - 2Rb - 2bc + ab).$$

Case V. $R > a + b$; intersection at L between AD , BC produced.


$$\text{Area} = \frac{1}{2}\pi R^2 + \frac{1}{2}\pi(R-c)^2 + \frac{1}{2}\pi(R-b+c)^2 + \text{sector } DTL + \text{sector } CIL + \text{triangle } DLC.$$

$$DL = R - a - c, DC = b, CL = (R - a - b + c).$$

$$\therefore \angle LDC = \theta = \cos^{-1} \left(\frac{b^2 + (R-a-c)^2 - (R-a-b+c)^2}{2b(R-a-c)} \right)$$

$$\angle DCL = \varphi = \cos^{-1} \left(\frac{b^2 + (R-a-b+c)^2 - (R-a-c)^2}{2b(R-a-b+c)} \right).$$

$$\therefore \text{Area} = \frac{1}{2}\pi(4R^2 + 2c^2 + b^2 - 2Rb - 2bc) + \frac{1}{2}(R-a-c)^2(\frac{1}{2}\pi - \theta) + \frac{1}{2}(R-a-b+c)^2(\frac{1}{2}\pi - \varphi) + \frac{1}{2}b(R-a-c)\sin\theta.$$



Case VI. $R > a + b$; intersection in angle KBM .

Area = $\frac{1}{2}\pi R^2 + \frac{1}{4}\pi(R-c)^2 + \text{sector } NDG + \text{sector } FBG + \text{triangle } CDG + \text{triangle } CBG$.

$DG = (R - a - c)$, $DC = b$, $BC = a$, $BG = (R - b + c)$, $DB = \sqrt{(a^2 + b^2)}$.

$$\therefore \angle DBG = \beta = \cos^{-1} \left(\frac{a^2 + b^2 + (R - b + c)^2 - (R - a - c)^2}{2(R - b + c)\sqrt{(a^2 + b^2)}} \right)$$

$$\angle GDB = \gamma = \cos^{-1} \left(\frac{a^2 + b^2 + (R - a - c)^2 - (R - b + c)^2}{2\sqrt{(a^2 + b^2)}(R - a - c)} \right)$$

$$\angle CBD = \beta' = \cos^{-1} \left(\frac{a}{\sqrt{(a^2 + b^2)}} \right) = \tan^{-1} \left(\frac{b}{a} \right)$$

$\angle CDB = \frac{1}{2}\pi - \beta'$, $\angle GDC = \gamma + \beta' - \frac{1}{2}\pi$, $\angle CBG = \beta - \beta'$, triangle $DCG + \text{triangle } BCG = \text{triangle } DGB - \text{triangle } DCB$.

$\therefore \text{Triangle } DCG + \text{triangle } BCG = \frac{1}{2}\sqrt{(a^2 + b^2)}(R - a - c)\sin\gamma - \frac{1}{2}ab$.

$\therefore \text{Area} = \frac{1}{4}\pi(3R^2 - 2Rc + c^2) + \frac{1}{2}(R - a - c)^2(\pi - \gamma - \beta') + \frac{1}{2}(R - b + c)^2(\beta - \beta') + \frac{1}{2}\sqrt{(a^2 + b^2)}(R - a - c)\sin\gamma - \frac{1}{2}ab$.

II. Solution by C. C. BEBOUT, Professor of Mathematics in Elgin High School, Elgin, Ill.; CHARLES C. CROSS, Libertytown, Md., and ELMER SCHUYLER, High Bridge, N. J.

As shown by the figure, the area grazed over is made up of one semi-circle, two quadrants, two sectors, and a triangle. We must sum the areas of these.

Area of semi-circle $HPK = \frac{1}{2}\pi R^2 \dots\dots(1)$.

Area of quadrant $HAK = \frac{1}{4}\pi(R - c)^2 \dots\dots(2)$.

Area of quadrant $KBI = \frac{1}{4}\pi(R - b + c)^2 \dots\dots(3)$.

By using formula $A = \sqrt{s(s-a)(s-b)(s-c)}$ we get area of triangle $DLC = \sqrt{[c(R-a)(R-a-b)(b-c)]} \dots\dots(4)$.

To find the areas of the sectors we must find the angles TDL and ICL . These angles are respectively the complements of angles LDC and DCL , which may be found as angles of the triangle LCD , whose sides are known. To find these angles use the trigonometric formula:

$$\tan \frac{1}{2}A = + \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$

$$\text{Then } \tan \frac{1}{2}\angle LDC = \sqrt{\frac{c(R-a-b)}{(R-a)(b-c)}} \text{ and } \tan \frac{1}{2}\angle LCD = \sqrt{\frac{(R-a-b)(b-c)}{c(R-a)}}$$

$$\therefore \angle LDT = (90^\circ - \angle LDC) = 90^\circ - 2\tan^{-1} \sqrt{\frac{c(R-a-b)}{(R-a)(b-c)}}$$

$$\text{and } \angle LCI = (90^\circ - \angle LCD) = 90^\circ - 2 \tan^{-1} \sqrt{\frac{(R-a-b)(b-c)}{c(R-a)}}.$$

$$\therefore \text{Area of sector } LDT = \frac{360^\circ}{90^\circ - 2 \tan^{-1} \sqrt{\frac{c(R-a-b)}{(R-a)(b-c)}}} [\pi(R-a-c)^2] \dots \dots (5),$$

$$\text{and area of sector } LCI = \frac{360^\circ}{90^\circ - 2 \tan^{-1} \sqrt{\frac{(R-a-b)(b-c)}{c(R-a)}}} [\pi(R-a-b+c)^2] \dots (6).$$

Summing (1), (2), (3), (4), (5), and (6), we get the area grazed over,

$$\begin{aligned} & \frac{1}{2} \pi (4R^2 + b^2 + 2c^2 - 2bR - 2bc) + \sqrt{[c(R-a)(R-a-b)(b-c)]} \\ & + \frac{360^\circ}{90^\circ - \tan^{-1} \sqrt{\frac{c(R-a-b)}{(R-a)(b-c)}}} [\pi(R-a-c)^2] \\ & + \frac{360^\circ}{90^\circ - 2 \tan^{-1} \sqrt{\frac{(R-a-b)(b-c)}{c(R-a)}}} [\pi(R-a-b+c)^2]. \end{aligned}$$

Solutions of problem 106 were received from P. S. Berg and Elmer Schuyler, and of problem 105 from Sylvester Robins. These solutions came too late for credit in last issue.

NOTE ON THE CALCULATION OF INTEREST AND DISCOUNT.

BY JOSEPH V. COLLINS, PH. D., PROFESSOR OF MATHEMATICS, STATE NORMAL SCHOOL, STEVENS POINT, WIS.

In the January number of the MONTHLY, Hon. J. H. Drummond, after giving the answer to the problem, 'What are the proceeds of a note discounted at a bank for 10 years at 10 per cent.,' as *nothing*, says, "The method of calculating discount used by banks was invented to evade the usury laws. I have thought that the court which first sustained the method could not have been well versed in mathematical principles."

A curious thing and commentary on the preceding is that all methods of calculating interest and discount are open to objection. Custom requires that interest be paid at the end of each year or specified fraction of a year. When it is not paid until the end of a period of two or more years, the lender is defrauded of the interest on his interest. If all interest payments are made promptly, it is equivalent to paying compound interest. The business world recognizes that compound interest is the only fair kind. Thus tables of bond values and the like are always made on a compound interest basis. But simple interest is